

# VACCINES NEEDED FOR SCHOOL

	All kindergartners and 4–6-year-old transfer students	All 7th graders and 7–18-year-old transfer students
Diphtheria, Tetanus, Pertussis (DTP, DTaP, Tdap) <sup>1</sup>	4 doses DTP or DTaP 1 dose must be at or after 4 years of age	4 doses diphtheria and tetanus or 3 doses if 1 <sup>st</sup> dose given on or after 1 year of age 1 dose Tdap at 11 years of age or older upon entry into 7 <sup>th</sup> grade or higher
Polio <sup>1</sup>	4 doses or 3 doses if dose 3 was given on or after 4 years of age	
Measles, Mumps, Rubella (MMR) <sup>1, 2</sup>	2 doses at or after 12 months of age	
Hepatitis B <sup>1, 2</sup>	3 doses	
Meningococcal Conjugate (MenACWY) <sup>1</sup>	None	1 dose at 11 years of age or older upon entry into 7 <sup>th</sup> grade or higher
Varicella (Chickenpox) <sup>1, 2</sup>	2 doses at or after 12 months of age or current lab immunity or history of varicella disease	

For more information please contact  
your healthcare provider, local  
pharmacist, or health department

[www.immunizationalliancemi.org](http://www.immunizationalliancemi.org)